

AD-A173 580


LASER-HEATING OF A TRANSPARENT CRYSTAL VIA ADSORBED
ATOMS(U) STATE UNIV OF NEW YORK AT BUFFALO DEPT OF
CHEMISTRY S VAN SMALEN ET AL OCT 86
UBUFFALO/DC/86/TR-15 N00014-86-K-0043


1/1

UNCLASSIFIED

F/G 20/13

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A173 580

OFFICE OF NAVAL RESEARCH
Contract N00014-86-K-0043
R & T Code 413f001---01
TECHNICAL REPORT No. 15

Laser-Heating of a Transparent Crystal Via Adsorbed Atoms

by

Sander van Smaalen, Henk F. Arnoldus and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

October 1986

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
OCT 20 1986
S D
D

DTIC FILE COPY

86 10 16 357

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/86/TR-15			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043	
8c. ADDRESS (City, State and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NOS.			
		PROGRAM ELEMENT NO.		PROJECT NO.	TASK NO.
					WORK UNIT NO.
11. TITLE Laser-Heating of a Transparent Crystal Via Adsorbed Atoms					
12. PERSONAL AUTHOR(S) Sander van Smaalen, Henk F. Arnoldus and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Yr., Mo., Day) October 1986	
				15. PAGE COUNT 19	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.	LASER HEATING ; PHOTON-PHONON CONVERSION;		
			TRANSPARENT CRYSTAL ; MASTER EQUATION;		
			VIA ADSORBED ATOMS ; PHOTODESORPTION		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A coated surface of a crystal is irradiated by intense infrared light. The optically active atomic bonds absorb photons from the laser field, and the subsequent spontaneous decay goes together with emissions of phonons into the crystal. This photon-phonon conversion results in an energy flux into the crystal. An equation for this flux is derived from the master equation for the level populations of the dressed atomic states. The saturation limit is discussed, and the general theory is illustrated with two examples. Furthermore, it is outlined with qualitative arguments that the quantum yield of photodesorption is not sensitive to the laser power. Key words					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE NUMBER (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

Physical Review B, in press

LASER-HEATING OF A TRANSPARENT CRYSTAL VIA ADSORBED ATOMS

Sander van Smaalen, Henk F. Arnoldus and Thomas F. George

Departments of Physics & Astronomy and Chemistry
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

ABSTRACT

A coated surface of a crystal is irradiated by intense infrared light. The optically active atomic bonds absorb photons from the laser field, and the subsequent spontaneous decay goes together with emissions of phonons into the crystal. This photon-phonon conversion results in an energy flux into the crystal. An equation for this flux is derived from the Master equation for the level populations of the dressed atomic states. The saturation limit is discussed, and the general theory is illustrated with two examples. Furthermore, it is outlined with qualitative arguments that the quantum yield of photodesorption is not sensitive to the laser power.

PACS: 71.36, 78.90



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

I. INTRODUCTION

The thermal desorption rate of atoms, which are bounded to the surface of a crystal and occupy a series of vibrational states, can be very low. Irradiation with resonant infrared laser light can enhance the desorption considerably,¹⁻⁶ especially at low temperatures. Then the thermal relaxation (coupling to the phonon field of the crystal) will tend to confine the atoms to the low-lying non-desorptive vibrational levels, but the laser will sustain a continuous excitation of the system. The balance between the thermal decay and the coherent excitation determines the level populations. Since atoms in high-lying states can more easily desorb into a continuum state through a resonant one-phonon transition than atoms in low-lying states, the driving laser will increase the desorption rate. The effect will be most pronounced as the optical frequency ω_L is close to an atomic-bond transition. However, the desorption rate as a function of ω_L is not tantamount to the optical absorption profile, due to the competing thermal relaxation. Indeed, an excited bond might decay to a lower state together with an emission of a phonon into the crystal. This process can be considered as a photon-phonon conversion, with a heat-flow into the crystal as a net result.^{7,8} Hence, the quantum yield of photodesorption will be diminished strongly by thermal relaxation. Experimental values acquire an order of magnitude of at most 10^{-2} , which implies that more than 99% of the absorbed energy ends up in the phonon field.² Conversely, for a crystal which is transparent at frequency ω_L , a dye coating on the surface can provide a medium for an efficient laser-heating of the solid.

In this paper, we derive a microscopic equation for the energy flux into the crystal. It contains the laser power and frequency, the temperature of the crystal, and the transition matrix elements of the atomic

potential well as parameters. The results can be applied as the source terms for the macroscopic diffusion equation, which describes the temperature distribution in the crystal.^{9,10} Our work extends earlier treatments^{7,8} in that we allow the irradiance to be arbitrarily intense, which will enable us to study, for instance, saturation effects.

II. POPULATIONS OF DRESSED STATES

The vibrational eigenstates of the atomic bond will be denoted by $|k\rangle$, their populations by $n_k(t)$, and the energy eigenvalues by $\hbar\omega_k$. The strong incident radiation will be assumed to couple only resonantly a ground state $|g\rangle$ and an excited state $|e\rangle$, with $\omega_e - \omega_g = \omega_0 > 0$. We indicate the detuning with the driving field by $\Delta = \omega_L - \omega_0$, and the coupling strength is expressed in the Rabi frequency $\Omega = |\underline{E}_0 \cdot \underline{\mu}|/\hbar$. Here, \underline{E}_0 is the amplitude of the electric component of the incident field, and $\underline{\mu}$ is the electric dipole moment of the transition. The parameter Ω^2 will be referred to as the laser intensity, since it is proportional to E_0^2 . The diagonalization of the Hamiltonian of the driven atomic bond is easily accomplished, and it amounts to a transformation of the bare states $|e\rangle$, $|g\rangle$ (eigenstates for $\Omega^2 = 0$) into the dressed states $|+\rangle$, $|-\rangle$.¹¹ This is illustrated in Fig. 1. The eigenvalues are

$$\hat{\omega}_{\pm} = \frac{1}{2}(\omega_e + \omega_g + \omega_L) \pm \frac{1}{2}\text{sgn}(\Delta)\sqrt{\Omega^2 + \Delta^2}, \quad (2.1)$$

with $\text{sgn}(\Delta)$ representing the sign of Δ . We indicate by $\hbar\hat{\omega}_k$ an eigenvalue of the dressed state $|k\rangle$.

The interaction with the phonon field of the crystal can now be considered as a coupling between dressed states.¹² We suppose that the separations between adjacent levels are smaller than the Debye frequency ω_D ,

which implies that we can omit multiphonon processes. The coupling strengths are governed by the matrix elements $\langle k|S|l\rangle$ of the atomic operator $S = dV/dz$, which is the derivative of the potential well in the direction perpendicular to the surface. In a previous paper¹¹ we derived an equation for the populations of the dressed states, as they are determined by the single-phonon relaxation constants a_{kl} . With a_{kl} the inverse lifetime of the transition $|k\rangle \rightarrow |l\rangle$, this Master equation reads

$$\frac{d}{dt}n_k(t) = \sum_l (n_l(t)a_{lk} - n_k(t)a_{kl}). \quad (2.2)$$

Here the summation runs over the dressed states $|l\rangle$ and the bare states $|k\rangle$, with $k \neq e, g$, since for $k = e, g$ a dressed state is identical to a bare state. In the derivation of Eq. (2.2) we adopted the secular approximation, which asserts that the level separations, and especially $|\Omega'|$, are sufficiently larger than the damping constants. Hence our approach applies to situations where the detuning or the laser power Ω^2 is large. This case can be considered to be complementary to the weak-field limit of Refs. 7 and 8.

In the expressions for the relaxation constants a_{kl} , we can separate the dependence on the optical parameters Ω and Δ from the pure thermal part. It appears that Ω and Δ only enter the expressions for the lifetimes in the combination g_{\pm} , defined by

$$g_{\pm} = \frac{1}{2} \{ 1 \mp (1 + \Omega^2/\Delta^2)^{-1/2} \}. \quad (2.3)$$

Hence, the distribution of population over the dressed states is governed by the single dimensionless optical parameter Ω^2/Δ^2 . In the strong-field limit $\Omega^2/\Delta^2 \rightarrow \infty$, both g_{+} and g_{-} tend to the finite asymptotic value of $\frac{1}{2}$, which indicates saturation.

The temperature-dependent part of the relaxation constants can be expressed in the field-free inverse lifetimes for transitions between bare states. In a simple Debye model the rate constants attain the form

$$a_{kl} = \frac{v'}{2\pi\hbar Mc^3} |\langle k|S|l \rangle|^2 \frac{\omega}{1 - \exp(-\hbar\omega/k_B T)} \Theta(\omega_D - |\omega|), \quad (2.4)$$

where v' is the volume of a unit cell, M is the mass of a crystal atom, c is the speed of sound, k_B is Boltzmann's constant, T is the temperature of the solid, Θ is the Heaviside step function, and $\omega = \omega_k - \omega_l$ is the level separation whose absolute value equals the phonon energy in a transition. The genuine impact of Eq. (2.4) on the thermal redistribution of level populations originates from its dependence on the sign of ω . A positive value of ω implies $\omega_k > \omega_l$, and hence the transition $|k\rangle \rightarrow |l\rangle$ is a decay of the system, whereas $\omega < 0$ reflects a thermal excitation of the same transition. For any two levels $|k\rangle, |l\rangle$, with $\omega_k > \omega_l$, both phonon absorption (excitation, a_{lk}) and phonon emission into the crystal (decay, a_{kl}) occurs, and it follows from Eq. (2.4) that $a_{kl} \gg a_{lk}$.

With these preliminary remarks, we can now write down the inverse lifetimes. If both levels $|k\rangle$ and $|l\rangle$ are each unequal to $|+\rangle$ or $|-\rangle$, the rate constant is simply a_{kl} from Eq. (2.4). For laser-assisted transitions we have¹¹

$$a_{\pm k} = g_{+}^2 a_{ek} + g_{\pm}^2 a_{gk}, \quad (2.5)$$

$$a_{k\pm} = g_{+}^2 a_{ke} + g_{\pm}^2 a_{kg}, \quad (2.6)$$

for $k \neq \pm$. The pure dressed-state transitions are governed by the four rate constants

$$a_{\pm\pm} = g_{+}^2 a_{ge} + g_{\pm}^2 a_{eg}, \quad (2.7)$$

$$a_{\pm\pm} = g_+g_-(a_{ge} + a_{eg}) . \quad (2.8)$$

A transition $|+\rangle \rightarrow |+\rangle$ or $|-\rangle \rightarrow |-\rangle$ does obviously not alter the populations of a dressed state. In the Master equation (2.2), the a_{kk} -contributions vanish indeed, but we will see in the next section that the $|+\rangle \rightarrow |+\rangle$ and $|-\rangle \rightarrow |-\rangle$ transitions do contribute to the energy transportation, as can be anticipated from Fig. 1. Therefore, we cannot discard these transitions in general. Furthermore, we note that a rate constant gains a contribution from two distinct processes, whenever a $|\pm\rangle$ state is involved. This is again clear from Fig. 1.

With Eqs. (2.3)-(2.8) the explicit dependence of the coefficients a_{kl} on the laser parameters, the temperature and the properties of the potential well is tracked down completely. For any configuration of states, the Master equation (2.2), accompanied by the normalization

$$\sum_k n_k(t) = 1, \quad (2.9)$$

can be solved immediately.

III. ENERGY FLUX

From the Master equation it follows that a transition $|k\rangle \rightarrow |l\rangle$ occurs at a rate $n_k(t)a_{kl}$ (number of transitions per unit of time). Every decay/excitation corresponds to an absorption/emission of a phonon by the crystal, with an energy equal to the level distance $\hbar(\hat{\omega}_k - \hat{\omega}_l)$. If we write $\hbar(\hat{\omega}_k - \hat{\omega}_l)$ for the gain of energy of the crystal in a single $|k\rangle \rightarrow |l\rangle$ transition, then the distinction between absorption and emission is included in the sign of $\hat{\omega}_k - \hat{\omega}_l$. Care should be exercised in the transitions to and

from the dressed states $|\pm\rangle$, since these transitions involve two diagrams with different phonon energy. Therefore, these rates should be divided accordingly, and for the gain of energy the contributions from the two doublets in Fig. 1 should be distinguished. It is immediately clear which part of the rate constant corresponds to a specific transition, since the argument $\omega_k - \omega_l$ in a_{kl} equals the phonon frequency. These notions yield the formal expression for the energy flux into the crystal (absorbed energy per unit of time)

$$\frac{dW}{dt} = \sum_{kl} \hbar(\hat{\omega}_k - \hat{\omega}_l) n_k(t) a_{kl}, \quad (3.1)$$

where the summation runs over all diagrams of Fig. 1, and $\hbar(\hat{\omega}_k - \hat{\omega}_l)$ is the appropriate phonon energy.

The double sum is easily evaluated after substitution of the explicit expressions (2.4)-(2.8) for the relaxation constants, Eq. (2.1) for the energies of the dressed states, and after application of the Master equation. We obtain

$$\begin{aligned} \frac{dW}{dt} = & - \sum_k \hbar \hat{\omega}_k \frac{d}{dt} n_k(t) \\ & + \hbar \omega_L \sum_{k \neq \pm} \{ a_{kg} n_k(t) - a_{gk} (g_+ n_+(t) + g_- n_-(t)) \} \\ & + \hbar \omega_L \{ a_{eg} (g_- n_+(t) + g_+ n_-(t)) - a_{ge} (g_+ n_+(t) + g_- n_-(t)) \}. \end{aligned} \quad (3.2)$$

Here the summations run over the bare states $k \neq e, g$ and over \pm , but no longer over diagrams. The right-hand side of Eq. (3.2) contains the bare-states inverse lifetimes a_{kl} , the optical parameters g_{\pm} and the populations $n_k(t)$ of the dressed states. This implies that dW/dt is expressed entirely in known parameters and in the solution $n_k(t)$ of the Master equation. The expressions in curly brackets are combinations of gain and loss terms

pertaining to the same transition, which is reminiscent of the structure of the Master equation.

IV. STEADY STATE

After a transient time of the order of a_{kl}^{-1} , the system will approach a steady state, due to the phonon relaxation. The populations tend to their long-time limit

$$\bar{n}_k = \lim_{t \rightarrow \infty} n_k(t), \quad (4.1)$$

and the Master equation reduces to

$$\sum_l \bar{n}_l a_{lk} = \sum_l \bar{n}_k a_{kl}. \quad (4.2)$$

In the energy equation (3.2), the first term on the right-hand side disappears, and the energy-flux dW/dt acquires a time-independent value in this limit.

In order to reveal more clearly the structure of the steady-state energy flux, we transform Eq. (3.2) to its bare-state equivalent. Due to the fact that the coherences with respect to the dressed states vanish in the long-time limit, the populations of the excited state and the ground state can be expressed in \bar{n}_+ and \bar{n}_- according to

$$\bar{n}_e = g_- \bar{n}_+ + g_+ \bar{n}_-, \quad (4.3)$$

$$\bar{n}_g = g_+ \bar{n}_+ + g_- \bar{n}_-. \quad (4.4)$$

Then Eq. (3.2) for $t \rightarrow \infty$ assumes the transparent form

$$\frac{dW}{dt} = \gamma_{\omega_L} \sum_{k=\text{bare}} (\bar{n}_k a_{kg} - \bar{n}_g a_{gk}). \quad (4.5)$$

Notice that the optical parameters g_{\pm} have vanished completely in this representation. They only enter the energy flux through the populations \bar{n}_k

of the states, which are affected by the laser field. The summation in Eq. (4.5) is a gain/loss balance for the population of the ground state $|g\rangle$. The first term (gain) reflects that a photon-absorption rate is maintained by a permanent driving of the transition $|g\rangle \rightarrow |e\rangle$ by the laser. The subsequent thermal relaxation to the ground state is accompanied with phonon emissions into the crystal, which yields the net energy flux. Conversely, the second term (loss) corresponds to stimulated photon emissions in $|e\rangle \rightarrow |g\rangle$ transitions. Then the crystal should provide the energy for the subsequent thermal redistribution.

The summation over all bare states in Eq. (4.5) can be eliminated with the Master equation (4.2) and the relations (4.3), (4.4) between bare-state and dressed-state populations. Recall that Eq. (4.2) pertains to dressed states, whereas the summation in Eq. (4.5) runs over bare states. The result can be cast in the remarkably simple form

$$\frac{dW}{dt} = \gamma_{\omega} \frac{\Omega^2}{L_{4\Delta}^2} (A_g + A_e)(\bar{n}_g - \bar{n}_e). \quad (4.6)$$

The optical prefactor arises from the combination

$$\frac{g_- g_+}{(g_- - g_+)^2} = \frac{\Omega^2}{4\Delta^2}, \quad (4.7)$$

and the total inverse lifetime A_k of a state $|k\rangle$ is defined as

$$A_k = \sum_{l=\text{bare}} a_{kl}. \quad (4.8)$$

The photon-absorption (-emission) rate is proportional to $\bar{n}_g(\bar{n}_e)$, and it appears that just the difference $\bar{n}_g - \bar{n}_e$ enters the expression for dW/dt . This is reminiscent of the standard result for the low-intensity absorption profile, but the expression (4.6) holds for arbitrary intensity Ω^2 .

V. SATURATION

It might appear from Eq. (4.6) that the energy flux is proportional to Ω^2 , and therefore can increase unlimited with the laser intensity. Such is however not the case, since for Ω^2/Δ^2 large, the population inversion $\bar{n}_e - \bar{n}_g$ tends to zero. In this section we shall derive an upper limit for dW/dt .

To this end, we first write Eq. (4.5) in the form

$$\frac{dW}{dt} = \hbar\omega_L \left\{ -\Lambda_g \bar{n}_g + \sum_{k=\text{bare}} \bar{n}_k a_{kg} \right\}. \quad (5.1)$$

From the Master equation and Eqs. (4.3) and (4.4), it is easy to derive the identity

$$\sum_{k=\text{bare}} \bar{n}_k (a_{kg} + a_{ke}) = \Lambda_g \bar{n}_g + \Lambda_e \bar{n}_e. \quad (5.2)$$

Expression (5.1) for dW/dt refers to transitions from and to the ground state, but with Eq. (5.2) we can convert the reference level to the excited state. This yields

$$\frac{dW}{dt} = \hbar\omega_L \left\{ \Lambda_e \bar{n}_e - \sum_{k=\text{bare}} \bar{n}_k a_{ke} \right\}, \quad (5.3)$$

and we note the complementary nature of Eqs. (5.1) and (5.3). Since $\bar{n}_k a_{ke} > 0$, we obtain from Eq. (5.3) the upper limit to the energy flux

$$\frac{dW}{dt} \leq \hbar\omega_L \Lambda_e \bar{n}_e. \quad (5.4)$$

With $\bar{n}_e \leq 1$, this reduces to $dW/dt \leq \hbar\omega_L \Lambda_e$, where the right-hand side is independent of the optical parameters. Hence, the inequality (5.4) implies that for strong fields the e,g transition saturates, and thus the energy flux is bounded. The limit (5.4), however, is an upper bound and not the saturation limit in general. The behavior of dW/dt for $\Omega^2/\Delta^2 \rightarrow \infty$ will depend on the phonon-relaxation parameters a_{kl} and the level configuration.

The inequality (5.4) with $\bar{n}_e = 1$ expresses the upper bound in the loss coefficient Λ_e of level $|e\rangle$. The complementary relation follows from Eq.

(5.1) and becomes

$$\frac{dW}{dt} \leq \hbar\omega_L \sum_k a_{kg}, \quad (5.5)$$

which pertains to the gain coefficient of the ground state.

VI. TWO LEVELS

Already the most simple case, the two-level system, exhibits an interesting feature. We discard the explicit transient solutions, since they only display the trivial exponential decay. The steady-state Master equation (4.2) is readily solved for \bar{n}_+ and \bar{n}_- . Then \bar{n}_e follows from Eq. (4.3), which gives

$$\bar{n}_e = \frac{(1-2g_+g_-)a_{ge} + g_+g_-(a_{eg}-a_{ge})}{(1-2g_+g_-)(a_{eg}+a_{ge})}, \quad (6.1)$$

and $\bar{n}_g = 1-\bar{n}_e$. With $A_e = a_{eg}$, $A_g = a_{ge}$, the energy flux becomes

$$\frac{dW}{dt} = \hbar\omega_L \frac{\Omega^2}{2\Omega^2 + 4\Delta^2} (a_{eg} - a_{ge}), \quad (6.2)$$

where we used the expression (2.3) for g_{\pm} . From the definition of the rate constants a_{kl} , Eq. (2.4), we deduce the properties

$$a_{kl}(T) - a_{lk}(T) = a_{kl}(T=0), \quad (6.3)$$

$$a_{lk}(T=0) = 0, \quad (6.4)$$

both for $\omega_k > \omega_l$. Combining everything yields the alternative formulation of Eq. (6.2) as

$$\frac{dW}{dt} = \hbar\omega_L \bar{n}_e(T=0) a_{eg}(T=0). \quad (6.5)$$

This reveals that the energy flux is independent of the temperature of the crystal, which is rather remarkable. From Eq. (6.5) we find the saturation limit to be

$$\frac{dW}{dt} = \frac{1}{2} \gamma \omega_L a_{eg}(T=0), \quad (6.6)$$

which is smaller than the upper-bound $\gamma \omega_L \bar{n}_e(T) a_{eg}(T)$ from Eq. (5.4).

Furthermore we notice that for $T = 0$ ($a_{ge} = 0$), the dependence of \bar{n}_e on the damping constant a_{eg} vanishes, so that the steady-state level populations are completely determined by the optical parameters Ω and Δ . This peculiar feature is a consequence of the assumed secular limit ($|\Omega'| \gg a_{eg}$).

VII. THREE LEVELS

Let us now consider the configuration with a third level $|\zeta\rangle$ present.

We then find

$$\begin{aligned} \frac{dW}{dt} = & \gamma \omega_L \frac{\Omega^2}{4\Delta^2} (A_e + A_g) ((a_{eg} - a_{ge}) A_\zeta + a_{e\zeta} a_{\zeta g} - a_{g\zeta} a_{\zeta e}) \\ & \times (a_{eg} A_\zeta + a_{g\zeta} A_e + a_{\zeta e} A_g + a_{e\zeta} a_{\zeta g} + a_{ge} a_{e\zeta} + a_{\zeta g} a_{ge} \\ & + \frac{\Omega^2}{4\Delta^2} (A_e + A_g) (a_{e\zeta} + a_{g\zeta} + 2A_\zeta))^{-1}, \end{aligned} \quad (7.1)$$

which becomes in the saturation limit $\Omega^2/\Delta^2 \rightarrow \infty$

$$\frac{dW}{dt} = \gamma \omega_L \frac{(a_{eg} - a_{ge}) A_\zeta + a_{e\zeta} a_{\zeta g} - a_{g\zeta} a_{\zeta e}}{a_{e\zeta} + a_{g\zeta} + 2A_\zeta}. \quad (7.2)$$

In the limit $A_\zeta^{-1} \rightarrow 0$, Eq. (7.1) reduces to the two-level result from the previous section, as it should.

In the limit $T \rightarrow 0$, the rates for upward transitions vanish. This implies that we have to distinguish three cases. First, for $\omega_\zeta > \omega_e > \omega_g$ we find again the results for the two-level case, since the population of $|\zeta\rangle$ becomes zero. For the situation $\omega_e > \omega_g > \omega_\zeta$ we obtain $dW/dt = 0$, as a result of the fact that all population resides in $|\zeta\rangle$. This prohibits any excitation by the laser field. The interesting case is $\omega_e > \omega_\zeta > \omega_g$, for which

we obtain

$$\frac{dW}{dt} = \gamma \omega_L A_e \frac{\Omega^2}{4\Delta^2 + \Omega^2(2 + a_{e\zeta}/a_{\zeta g})} . \quad (7.3)$$

This exhibits clearly that a non-zero value of the ratio $a_{e\zeta}/a_{\zeta g}$ diminishes the energy transfer to the crystal. More generally, this idea can be inferred from Eq. (5.3). Any appearance of additional levels with a finite population tends to reduce the energy flux. The population of the excited state for $T = 0$ and $\omega_e > \omega_\zeta > \omega_g$ is found to be

$$\bar{n}_e = \frac{\Omega^2}{4\Delta^2 + \Omega^2(2 + a_{e\zeta}/a_{\zeta g})} . \quad (7.4)$$

Comparison with Eq. (7.3) then yields

$$\frac{dW}{dt} = \gamma \omega_L A_e \bar{n}_e , \quad (7.5)$$

which is reminiscent of the result (6.5) for the two-level case and of the saturation constraint (5.4).

VIII. CONCLUSIONS

We have studied the surface-adsorbate mediated conversion of laser photons into phonons of a harmonic crystal. A net energy flux dW/dt into the crystal emerges from the different laser-assisted single-phonon transitions. We have derived a variety of expressions for dW/dt , which include the explicit dependence on the laser power, the detuning, the temperature and the properties of the vibrational bond. This was achieved by a proper interpretation of the thermal transition rates, as they appear in the Master equation with respect to the dressed states. Our approach applies to arbitrarily strong incident fields and to any configuration of atomic levels, but is restricted to one-phonon processes. We digressed on saturation effects to derive an upper limit for dW/dt . The general

treatment was exemplified by an explicit evaluation of dW/dt for the cases of a two-level and a three-level system. We find that for the two-level system the energy flux is independent of the temperature of the crystal. For three levels and $T \rightarrow 0$, we find the relation $dW/dt = \hbar\omega_L A_{eg} \bar{n}_e$, which can readily be understood. The only process which gives rise to energy transportation is an optical excitation $|g\rangle \rightarrow |e\rangle$ and a successive thermal decay $|e\rangle \rightarrow |g\rangle$ or $|e\rangle \rightarrow |\zeta\rangle \rightarrow |g\rangle$. The rate for this process equals $A_{eg} \bar{n}_e$, and the energy gain per transition is $\hbar\omega_L$. The effect of the laser is reflected in $\bar{n}_e \neq 0$, since for $\Omega^2 \rightarrow 0$ the system is bound to be in the ground state $|g\rangle$.

Another interesting feature can be deduced immediately from our results. If the laser is used for photodesorption of the atoms, then the desorption rate will roughly be proportional to the sum of the populations of the high-lying states. For the case of a three-level system with $\omega_e > \omega_\zeta > \omega_g$, the number of desorbed atoms per unit of time will be $\bar{n}_e a_{ec}$, where a_{ec} is the rate constant for a transition to the continuum. On the other hand, the number of absorbed photons per unit of time equals $(dW/dt)/\hbar\omega_L$. If we define the quantum yield as the ratio of these two numbers, then we find from Eq. (7.5) that it becomes a_{ec}/A_{eg} for low temperatures and in the three-level case, and in general for a two-level system. This ratio is independent of Ω^2/Δ^2 , so the efficiency of photodesorption cannot be enhanced by tuning the laser or increasing the power. The absolute rate is however proportional to \bar{n}_e , which depends on the intensity. For not too strong fields we have $\bar{n}_e \propto \Omega^2$, which is not surprising. Resonance effects appear if we tune ω_L into the line center of the absorption profile ($-\omega_0$). Then \bar{n}_e approaches its saturation limit ($\frac{1}{2}$), which provides the maximum photodesorption yield.

ACKNOWLEDGEMENTS


This research was supported by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009, the Office of Naval Research, and the National Science Foundation under Grant CHE-8519053. The United States Government is authorized to reproduce and distribute reprints notwithstanding any copyright notation hereon.

REFERENCES

1. M.S. Djidjoev, R.V. Khokhlov, A.V. Kiselev, V.I. Lygin, V.A. Namiot, A.I. Osipov, V.I. Pachenko and B.I. Provotorov, in Laser Chemistry at Surfaces, edited by A. Mooradian, T. Jaeger and P. Stokseth (Springer, Berlin, 1976), p. 100 ff.
2. J. Heidberg, H. Stein and E. Riehl, Phys. Rev. Lett. 49, 666 (1982).
3. T.J. Chuang and H. Seki, Phys. Rev. Lett. 49, 382 (1982).
4. T.J. Chuang, Surf. Sci. Rep. 3, 1 (1983).
5. I. Hussla, H. Seki, T.J. Chuang, Z.W. Gortel, H.J. Kreuzer and P. Piercy, Phys. Rev. B 32, 3489 (1985).
6. K. Veeken, P.A.M. Van Der Heide, L.M. ten Dam, A.R. de Vroomen and J. Reuss, Surf. Sci. 166, 1 (1986).
7. Z.W. Gortel, H.J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 28, 2119 (1983).
8. Z.W. Gortel, H.J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B 27, 5066 (1983).
9. J.T. Lin and T.F. George, J. Appl. Phys. 54, 382 (1983).
10. D. Burgess, Jr., P.C. Stair and E. Weitz, unpublished (1986).
11. H.F. Arnoldus, S. van Smaalen and T.F. George, Phys. Rev. B, submitted.
12. C. Cohen-Tannoudji and S. Reynaud, J. Phys. B: At. Mol. Phys. 10, 345 (1977).

FIGURE CAPTION

Fig. 1: The laser field couples the atomic levels $|e\rangle$ and $|g\rangle$. Bare-state transitions occur as radiationless or laser-assisted single-phonon excitations or decays. Diagonalization of the dipole interaction transforms the set $|e\rangle, |g\rangle$ into a ladder of dressed states $|i\rangle$, where the doublets are separated by the photon energy $\hbar\omega_L$. The distance between $|+\rangle$ and $|-\rangle$ in a single doublet equals $(\Delta^2 + \Omega^2)^{1/2}$, and the absolute position of the upper set of states is $\hat{\omega}_\pm$ from Eq. (2.1). The occurring single-phonon transitions are indicated. The state $|m\rangle$ couples in four ways with the ladder of dressed states, where each transition has a different phonon energy. There are four couplings between the two doublets, but $|i\rangle \rightarrow |i\rangle$ transitions in a single doublet are not present. The unperturbed states $|m\rangle$ and $|n\rangle$ couple only through a resonant one-phonon process. The transitions of $|n\rangle$ to the ladder of dressed states are suppressed in this picture.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
x Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Doua Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
x Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSROC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
x Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		x Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

X Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

END

11-86

DTIC